

VASCULAR MANIFESTATIONS OF BEHCET DISEASE


1^{er} Auteur : Oumayma, Albouchi, (1) Service de Médecine interne, Hopital Habib Thameur, Tunis, Tunisie;
Autres auteurs, équipe:I. Arbaoui (1) ; M. Somaï (1) ; R. Boukhzar (1) ; B. Ben Dhaou (1) ; F. Boussema (1) ; I. Rachdi (1) ; Z. Aydi (1) ; F. Daoud (1)
(1) Service de Médecine interne, Hopital Habib Thameur, Tunis, Tunisie;

Introduction: Vascular lesion of BD can occur in large, medium or small arteries or veins. Clinical features of vascular BD are various. Our study aimed to investigate characteristics of vascular BD in both genders in different age.

Patients and methods: This retrospective study included patients followed up between 2010 and 2022 for BD in an internal medicine department.

Number of patients	149
Incidence rate of vascular Behcet’s disease	30,87% (46/149)
Sex ratio	8 men 2 women
Age at diagnosis	34,7 (extremes of 20 and 59 years old)
Percentage as a revealing mode	22,14 (33/149)

TYPES AND LOCATIONS OF VENOUS THROMBOSIS



Conclusion: Our results are in accordance with the literature, which reports a prevalence of angio-Behçet estimated at between 22 and 25%, with a sex ratio of 5:1, an age range of between 25 and 40 years, and venous involvement accounting for almost 90% of vascular damage. The results of our study deserve to be elucidated by large-scale multicentric studies.

